

IAP20 Rec'd PCT/PTO 07 FEB 2006

**Fermented food product comprising soy protein****Description****5 Field of the invention**

The present invention relates to a food product fermented with lactic acid bacteria comprising soy protein. The invention further relates to use of such food products as a health food and a process for the preparation of the food  
10 products.

**Background to the invention**

According to the Codex Alimentarius (FAO/WHO 1977), yogurt is milk fermented with *Lactobacillus bulgaricus* and  
15 *Streptococcus thermophilus* and the microorganisms in the final product must be "viable and abundant". Other food products fermented with *S. thermophilus*, which may not be called yogurt according to the above definition exist. For instance "Bifidus yogurt" is made using selected cultures  
20 of *B. bifidum*, *B. longum*, *S. thermophilus*, and *L. delbrueckii*, which results in a yogurt-like product having a milder taste than yogurt.

Also known are so-called "soy yogurts", in which the main  
25 source of protein is not milk, but soy protein and which are fermented with yogurt culture. These soy yogurts are herein also described as yogurts. Examples of commercially available soy yogurts are Yofy produced by Van der Moortele and Sojasun produced by Triballat. Though the preparation  
30 of these commercial food products is not always known, most of these soy yogurts are prepared from soy beans and have a soy protein content that is relatively low.

US 6013771 describes isoflavone rich soy protein isolates and their application in food products. Column 8, Formulation 4, shows the preparation of a soy yogurt with SPI containing about 8 grams of soy protein per 170 gram 5 serving.

We have found that a food product prepared with soy protein isolate (SPI) as sole protein source is very thick and gives a non-dairy yogurt characteristic dry mouthfeel.  
10

We have found that the use of hydrolysed SPI in yogurt applications results in improved product stability at low pH, but the products have no yogurt texture. The texture/firmness in these products may be improved by  
15 addition of thickeners, such as starch or pectin. However, addition of these thickeners still does not create a dairy-like yogurt texture. Further for fruit yogurt applications additional problem is that the shear required to mix the fruit preparation into the yogurt further breaks down the  
20 texture.

#### Summary of the invention

It is therefore an object of the invention to provide a food product fermented with one or more lactic acid  
25 bacteria comprising soy protein that has a high level of soy protein and a good taste and texture resembling that of yogurt. It is a further object of the invention to provide such food products in which the presence of thickeners or gums is not required. Still a further object is to provide  
30 a food product that is a stirred-type food product, preferably spoonable as defined herein, in which flavouring

such as fruit can be easily mixed without disrupting the texture of the product.

One or more of these objects are attained according to the invention in that characterised in that the food product comprises at least 0.2 wt.% hydrolysed protein. According to the invention dairy-like spoonable yogurt texture may be obtained even without the need of thickeners.

#### 10 Detailed description of the invention

Lactic-acid Bacteria herein are gram-positive, rod- or sphere-shaped bacteria that produce lactic acid as the principal or sole end product of carbohydrate fermentation.

15 The amount of soy protein in the food product according to the invention is 4-10 wt.%, preferably 4-7 wt.%.

The amount of hydrolysed protein in the food product according to the invention is at least 0.2 wt., preferably, 20 at least 0.5 wt.% more preferably at least 1 wt.% and most preferably at least 1.5 wt.%. Amounts of hydrolysed soy protein present are included in the amounts of soy protein.

The presence and amounts of proteins (non-hydrolysed and 25 hydrolysed) present in the food product may be determined by the skilled person using common analysis techniques e.g. mass spectroscopy and liquid chromatography.

The production of the fermented soy according to the 30 invention may include the following steps:

- 1) Standardizing of soy starting material
- 2) Thermal treatment and homogenisation

- 3) Addition of starter culture
- 4) Fermentation at 40-45°C
- 5) Cooling to about 10°C and packaging
- 6) Mixing in fruit other flavouring components

5

The characteristic flavor of yogurt is largely due to carbonyl compounds, particularly acetaldehyde (25 - 50 mg/kg), volatile fatty acids, and to some extent, due to acetone and acetoin.

10

The typical pH of soy yogurt varies from 4.0 - 5.0, according to type of yogurt, fat content, and specific starter cultures used.

- 15 Low-acidity yogurts have a pH above 4.6, while normal acidity yogurt has a pH of 4.0-5.0.

The soy yogurt according to the invention may be set-style yogurt with a firm gelled texture, or may be stirred-style  
20 yogurt, with a spoonable or fluid texture. Fruit yogurts are preferably stirred type.

Preferably the fermentation for the set-style soy yogurt takes place in the package of the yogurt after packaging.

25

Stirred-type soy yogurt is preferably fermented in fermentation tanks. After fermentation, the product may be stirred at a low speed prior to its transport by pumping and filling. A moderate to low shear rate is critical to  
30 achieve the desired viscous properties of stirred-style yogurt.

The food products according to the invention may be frozen like ice cream or it may be dehydrated to give yogurt powder.

5 Non-hydrolysed soy protein isolate, when used alone in the food products results in very firm and dry textures at soy protein levels above 4 wt.%. Such food products also have a taste that is less preferred, in particular they give a dry mouthfeel.

10

Further texture improvement can be achieved by addition of some milk protein, e.g. skim milk powder (SMP) buttermilk powder (BMP) and/or whey powder (such as sweet whey powder), which will result in a smoother texture and better taste, in which case the soy protein isolate ratio still largely determines the texture.

Although firmness of the food products according to the invention is lower for all products after application of shear for instance for mixing in fruit preps, still spoonable textures may be achieved.

Preferably vegetable oil and/or may be added to the food products. The addition of oil leads to an improvement in terms of smoothness and appearance.

Although the addition of thickeners is not needed in food products according to the invention, thickeners may be added, e.g. for further improving syneresis (e.g. pectin, carrageenans, locust bean gum).

The food product may be fermented with any lactic acid bacteria or any combinations of lactic acid bacteria.

Preferably the food product is fermented with *Streptococcus*  
5 *thermophilus* and *Lactobacillus bulgaricus*. The combination of these lactic acid bacteria gives a characteristic yogurt taste.

The hydrolysed protein is preferably hydrolysed soy  
10 protein.

Preferably the food product has a Stevens value of 50 to 300 g, more preferably 50 to 250 g, even more preferably 120 to 200 g. Preferably the food product is spoonable.  
15

Preferably the food product has a viscosity of 10000 to 800000 mPa.s, more preferably 50000 to 200000, most preferably 50000 to 100000 mPa.s.

20 Preferably the food product comprises 0.1-3 wt.% milk protein.

The presence of oil and/or fat is advantageous in the products according to the invention. Oil and/or fat has a  
25 positive effect on mouthfeel, the products become more creamy, smoother and less watery. Additionally, the preferred addition of oil or fat make it easier to control foaming of the protein during preparation of the food product. Therefore preferably, the food product comprises  
30 0,2-10 wt.%, more preferably 0,5-5, even more preferably 0,5-2 wt.% oil or fat.

The invention further relates to a process for the preparation of a fermented food product, wherein a starting material comprising soy protein is fermented with on or more lactic acid bacteria and wherein 40 wt.% or more 5 relative to the total amount of soy protein is hydrolysed soy protein.

The starting material may comprise a partially hydrolyzed soy protein or it may alternatively comprise a mixture of 10 non-hydrolysed and hydrolysed protein.

Preferably, the starting material comprises a mixture of non-hydrolysed soy protein and hydrolysed soy protein. More preferably the mixture comprises 50-95 wt.% of hydrolysed 15 soy protein relative to the total amount of soy protein.

Preferably the mixture is prepared from hydrolysed soy protein isolate (herein also designated as SPI) and non-hydrolysed soy protein isolate.

20

### Examples

#### Stevens value

Stevens values give an indication about the firmness 25 (spoonability) of a product. The firmness of all products at 5°C (after storage at 5°C for 24 hours) was measured using a Stevens Texture Analyser (1 mm/sec, 25 mm depth, mayonnaise grid (mesh 7, thread thickness 0.8 mm, mesh width 2.8 mm) and is quoted herein as the Stevens value (in 30 g). The accuracy of this measurement in all cases is  $\pm 10$  g.



Stevens values of above 120 and well below 300 g are typical for a spoonable product.

#### Viscosity

- 5 Viscosity of the food products is measured using a Brookfield viscometer, spindle 6, at 5 rotations per minute (herein abbreviated rpm).

#### Syneresis

- 10 Syneresis was measured by after storing flasks with the food product for 24 hours at 25°C and subsequently 24 hours at 5°C. The weight of the flasks before ( $w_i$ ) and after ( $w_d$ ) decantation of supernatant water and syneresis values were expressed as  $(w_i - w_d) / w_i$ .

15

#### Bostwick

- The Bostwick equipment consists of a 125 ml reservoir provided with a outlet near the bottom of a horizontally placed rectangular tub and closed with a vertical barrier.
- 20 The tub's bottom is provided with a 25 cm measuring scale, extending from the outlet of the reservoir. When equipment and sample both have a temperature of 5°C, the reservoir is filled with 125 ml of the sample after it has been shaken by hand ten times up and down. When the closure of the
- 25 reservoir is removed the sample flows from the reservoir and spreads over the tub bottom. The path length of the flow is measured after 30 seconds. The value, expressed as cm per 30 seconds is the Bostwick ratio.

30



**Example 1**

A fermented food product (yogurt) was prepared as follows. Soy protein isolate Supro Plus<sup>TM</sup> 651 and hydrolysed soy protein isolate FXP H220D, both containing about 80 wt.% soy protein and originating from Protein Technology International, skimmed milk powder and salt in amounts as given in table 1 were mixed with hot demi water at 70°C using a Silverston mixer at a speed of 2500 rpm until a homogeneous mixture is obtained. The mixture was sterilized in an Ultra High Temperature (UHT) apparatus at 170-200 bar, with heating bath temperature 85°C and holding bath temperature 75°C. The sterilized product was filled into 125 ml flasks and 0.2 wt.% yogurt culture YF202 (ex Rhodia) suspended in tripton solution was added to the flasks. The flasks were incubated at 43°C for 12 hours and thereafter cooled to 10°C and stored at 5°C.

Viscosity, hardness and syneresis of the prepared yogurt-type food compositions were measured according to the above described test methods. The results are given in table 2.

**Examples 2-4 and comparative experiments A and B**

The food products in these experiments were prepared as in example 1, however with different amounts of non-hydrolysed and hydrolysed soy protein isolates as indicated in table 1.

The results are given in table 1.

Examples 1-4 and the comparative experiments show that when only non-hydrolysed soy protein isolate is used, too thick products are prepared that have a less preferable taste (mouthdry). When only hydrolysed soy protein isolate is used no spoonable yogurt texture may be obtained.

The yogurts of examples 1-4 had a clean, fresh yogurt taste with no or little residual soy taste

10 Fruit yogurts may be made by adding fruit or fruit puree to the yogurts of examples 1-4.

Table 1: Composition of milk starting material

Composition	Example 1 (wt.%)	Example 2 (wt.%)	Example 3 (wt.%)	Example 4 (wt.%)	Comp. Ex. A (wt.%)	Comp. Ex. B (wt.%)
Ratio	70/30	60/40	40/60	20/80	100/0	0/100
hydrolysed/ non-hydrolysed soy						
Hydrolysed SPI FXP H0220D	5.84	5.00	3.13	1.56	7.81	0.00
Non-hydrolysed SPI Supro 651	2.47	3.29	4.63	6.17	0.00	8.23
Sunflower oil	1.50	1.50	1.50	1.50	1.50	1.50
Salt (NaCl)	0.10	0.10	0.10	0.10	0.10	0.10
Sweet milk powder	3.00	3.00	3.00	3.00	3.00	3.00
Demi water	87.09	87.11	87.64	87.67	87.59	87.17

Table 2: Properties of fermented food products, Bostwick values "—" means too low to be measured.

Example	Ratio hydrol./non- hydrol.SPI	pH	Viscosity (mPa.s)	Stevens value (g)	Bostwick value	Syneresis after 24 hr	Viscosity after shear (mPa.s)	Stevens value after shear (g)	Bostwick value after shear
A	100/0	4.46	70000	44	9	1.62	11200	12	10.5
1	70/30	4.51	278000	184	1.9	1.25	64000	52	2.5
2	60/40	4.49	320000	244	0.75	1.13	84000	62	2.2
3	40/60	4.45	295000	361	--	1.08	109000	105	1.8
4	20/80	4.46	774000	590	--	0.63	225000	197	--
B	0/100	4.49	770000	1111	--	0.85	386000	814	--

### Claims

1. Food product fermented with one or more lactic acid bacteria, comprising 4-10 wt.% soy protein, characterised in that the food product comprises at least 0.2 wt.% hydrolysed protein.
2. Food product according to claim 1, wherein the food product comprises at least 0.5 wt.% hydrolysed protein.
3. Food product according to claim 2, wherein the food product comprises at least 1.5 wt.% hydrolysed protein.
4. Food product according to claim 3, wherein the food product is fermented with *Streptococcus thermophilus* and *Lactobacillus bulgaricus*.
5. Food product according to any of claims 1 to 4, wherein the hydrolysed protein is hydrolysed soy protein.
6. Food product according to any of claims 1-5, wherein the food product comprises 0.2-10 wt.% oil and/or fat.
7. Food product according to claims 6, comprising 0.5-2 wt.% oil and/or fat.
8. Food product according to any of claims 1 to 7, wherein the food product has a Stevens value of 50 to 300 g.
9. Food product according to any of claims 1 to 8, having a viscosity of 10000 to 800000 mPa.s.

10. Food product according to any of claims 1 to 9, wherein the food product comprises 0.1-3 wt.% milk protein.
11. Process for the preparation of a fermented food product, wherein a starting material comprising soy protein is fermented with one or more lactic acid bacteria, characterised in that 40 wt.% or more relative to the total amount of soy protein is hydrolysed soy protein.
12. Process according to claim 11, wherein the starting material comprises a mixture of non-hydrolysed soy protein and hydrolysed soy protein.
13. Process according to claim 12, wherein the mixture comprises 50-95 wt.% of hydrolysed soy protein relative to the total amount of soy protein.
14. Process according to claim 12 or 13, wherein the mixture is prepared from hydrolysed soy protein isolate and non-hydrolysed soy protein isolate.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/007610

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A23C9/13 A23J3/16

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A23C A23J A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, FSTA, BIOSIS, COMPENDEX

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 198425 Derwent Publications Ltd., London, GB; Class D13, AN 1984-155386 XP002260893 & JP 59 082050 A (FUJI OIL CO LTD) 11 May 1984 (1984-05-11) abstract	1-10
Y A		12-14 11
X	WO 03/055324 A (RAISIO BENECOL OY ;WESTER INGMAR (FI); KUUSISTO PAEIVI (FI)) 10 July 2003 (2003-07-10) claims 1,4,16-18; example 10	11
Y A		12-14 1-10
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

22 October 2004

Date of mailing of the international search report

02/11/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Koch, J



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/007610

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 199408 Derwent Publications Ltd., London, GB; Class D13, AN 1994-063488 XP002260894 & SU 1 789 172 A (LENGD REFRIG IND TECHN INST) 23 January 1993 (1993-01-23)	11
A	abstract	1-10, 12-14
A	----- US 6 013 771 A (SPADAFORA FRANK E ET AL) 11 January 2000 (2000-01-11) cited in the application column 8, lines 1-29 -----	1-12

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/007610

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
JP 59082050	A	11-05-1984	JP	1629235 C	20-12-1991
			JP	2054058 B	20-11-1990
WO 03055324	A	10-07-2003	CA	2470367 A1	10-07-2003
			EP	1455588 A1	15-09-2004
			WO	03055324 A1	10-07-2003
SU 1789172	A	23-01-1993	SU	1789172 A1	23-01-1993
US 6013771	A	11-01-2000	CA	2307064 A1	28-10-2001

**This Page Blank (uspto)**